

# Agromet Advisory Bulletin for the Nagapattinam District

#### (From 04th October 2023 to 08th October 2023)



Date: 03.10.2023

# Issued jointly by Tamil Nadu Rice Research Institute - Aduthurai and IMD

**Bulletin no:** 79/2023

Weather Summary for the period since the issue of last bulletin no. 78 (29.09.2023) for Tamil Nadu: Nagapattinam district

Maximum temperature of 33 - 35 °C, minimum temperature of 24 - 27 °C, isolated rainfall was recorded in Nagapattinam district.

#### Weather Summary and forecast for Nagapattinam district

Past week Summary 29.09.2023 to 02.10.2023				Parameters	Ensemble weather forecast valid until 08.30 hr of 08.10.2023				
Day-1	Day-2	Day-3	Day-4	Date	Day-1	Day-2	Day-3	Day-4	Day-5
29/09	30/09	01/10	02/10		04/10	05/10	06/10	07/10	08/10
4.6	0.0	0.2	0.0	Rainfall (mm)	0	0	3	3	0
35	34	33	35	Max. Temp. (°C)	35	35	35	35	34
24	27	27	26	Min. Temp. (°C)	26	26	25	25	25
5	5	5	4	Cloud cover (Octa)	3	3	4	7	3
79-76	73-73	78-75	80-75	RH I-II (%)	80-50	80-50	90-50	80-50	80-60
15	10	11	13	Wind (kmph)	20	22	20	20	20
South	South South West	South	South	Wind direction	South South West	South South West	South South West	South South West	South South West

#### Weather forecast for next five days (04.10.2023 to 08.10.2023)

According to district level forecast issued by India Meteorological Department for Nagapattinam district, the sky will be clear. Light rainfall is expected on 06.10.2023 and 07.10.2023. Maximum temperature is expected to be around 34 - 35 °C and minimum temperature is expected to be around 25 - 26 °C. Morning relative humidity is expected to be around 80 - 90 percent and evening relative humidity is expected to be around 50 - 60 percent. Average wind speed is expected to be around 20 - 22 km per hour. The wind direction will be south south west.

## **Agro Advisory**

Crop	Stage	Advisory			
Ger	neral	Farmers are requested to download and utilize the agricultural based mobile apps suggested by TNAU (TNAU AAS, Meghdoot and DAMINI).			
Rice	Main field	By utilizing dry weather, fertilizers and pesticides can be applied.  Apply 32.6 kg Urea and 8.4 kg of MOP should be applied after 35 days after transplanting for 1 acre rice field.  The prevailing weather condition favors Thrips infestation. To control the Thrips damage in transplanted rice, apply Thiamethxam 25WG @ 40g/ acre.			
cotton		Due to prevailing weather condition there are chances of cotton green leaf hopper infestation. To control spray thiamethxam 25 WG 0.6 gram/litre or imidacloprid 17.8 SL 0.6 ml/litre			
Cattle	Cow	Before monsoon ensure that cows and buffaloes were vaccinated against food and mouth disease.			

## **SMS advisory**

Farmers are requested to download and utilize the agricultural based mobile apps suggested by TNAU (TNAU AAS, Meghdoot and DAMINI).

Director TRRI, Aduthurai