

Pandit Jawaharlal Nehru College of Agriculture and Research Institute, Karaikal - 609 603

Agromet Advisory Bulletin (AAB) for KARAIKAL District Issued jointly by IMD & PAJANCOA & RI



Phone No. 04368 – 261182 E-mail: <u>agronaas2012@gmail.com</u>

Fax: 04368-261260

Bulletin No. 029 (04)/PJN/GKMS/2024

Weather Summary for Karaikal district

Past week Summary (04.04.2024 to 08.04.2024)					Weather Parameters	Ensemble weather forecast valid until 08.30 hrs of 14.04.2024				
04/04	05/04	06/04	07/04	08/04		Day-1 10/04	Day-2 11/04	Day-3 12/04	Day-4 13/04	Day-5 14/04
0	0	0	0	0	Rainfall (mm)	0	0	2	2	0
36	37	39	39	37	Max. Temp. (°C)	35	34	34	34	34
27	26	25	25	26	Mini. Temp. (°C)	27	27	26	27	26
2	2	2	1	4	Cloud cover (Octa)	3	4	6	5	6
60-87	57-92	48-89	58-7	56-92	RH Even- Morn. (%)	50-70	50-80	60-80	50-80	50-80
6	4	4	4	4	Wind (kmph)	16	16	16	18	18
WNW	WNW	WNW	WNW	SSW	Wind direction	Е	ESE	ESE	ESE	ESE

Forecast for next five days: 10.04.2024 to 14.04.2024

- During next five days the sky will be partly cloudy and light drizzle may occur on 12th and 13th April. The temperature range is expected to be around 26-35°C. Relative humidity range is expected to be around 50-80%. The wind speed is expected to be 16-18 km per hour and the wind will be mostly from east south east direction.
- * Rainfall (1.82 mm), maximum temperature (34.78°C) and minimum temperature (24.70°C) will be normal from 14th to 20th April.

Agro Advisory

Crop/livestock	Stage/Particulars	Agro Advisory					
Blackgram/	Flowering and pod development	To manage powdery mildew in blackgram spray Propiconazole @ 1 ml/l or Carbendazim @ 1 ml/l.					
Greengram	Leaf Crinkle	To manage leaf crinkle disease in green gram, control the white fly (vector of leaf crinkle) by spraying thiamethoxam 25 WG @ 0.2 g/l.					
Cotton	Vegetative growth	To manage weeds in cotton, apply post emergence herbicide pyrithiobac sodium @ 62.5g a.i./ha + quizalofop ethyl @ 50 g a.i./ha at 20-25 DAS.					
Brinjal	Phomopsis blight	To manage Phomopsis blight in brinjal, spray Pseudomonas fluorescens or Bacillus subtilis @ 5g or 5ml/1.					
Cattle	Babesiasis	In cattle, as a preventive measure for Babesiasis disease, apply Butox @ 2 ml/l of water on cattle and also spray it on the nearby areas of the cattle shed to control ectoparasites.					

 $\pmb{SMS}\text{:}$ Light drizzle may occur on 12^{th} and 13^{th} April.

Date: 09.04.2024