# Tamil Nadu Agricultural University



India Meteorological Department Agromet Advisory Bulletin for the Tiruppur District Bulletin No.017/2024 Issued on 27.02.2024 (From 28<sup>th</sup> February 2024 to 03<sup>rd</sup> March 2024) Issued jointly by Agro Climate Research Centre, TNAU and IMD



#### Past Weather for Tamil Nadu District

Weather Summary for the period since the issue of last bulletin 016 (23.02.2024) for Tamil Nadu. Maximum temperature ranged from 32°C to 36°C and minimum temperature ranged from 23°C to 27°C, over the State. Kodaikanal and Uthagamandalam recorded maximum temperature ranged from 18°C to 24°C and minimum temperature ranged from 10°C to 11°C. During this period isolated rainfall was recorded in Tamil Nadu.

#### **Tiruppur District Past Weather**

In Tiruppur district, during 23.02.2024 to 26.02.2024 the Maximum temperature was recorded was 34°C to 37°C and the minimum temperature was 22°C to 24°C. The morning relative humidity recorded was 83 per cent and evening relative humidity was 41 per cent. The average wind speed recorded was 8-16 km per hour and the wind direction were from East direction.

Past week Summary (23.02.2024 to 26.02.2024)				Ensemble weather forecast valid until 08.30 hrs of 03.03.2024						
Day-1 23/02	Day-2 24/02	Day-3 25/02	Day-4 26/02	Date	Day-1 28/02	Day-2 29/02	Day-3 01/03	Day-4 02/03	Day-5 03/03	
0.0	0.0	0.0	0.0	Rainfall	0	0	0	0	0	
36	37	36	34	Max. Temp. (° C)	34	34	35	35	36	
23	23	24	22	Mini. Temp. (° C)	23	24	24	25	25	
4	6	3	5	Cloud cover (Octa)	2	3	2	4	1	
80	83	80	78	RH morning (%)	80	70	80	70	70	
41	47	52	52	RH evening (%)	20	20	20	20	20	
9	8	9	16	Wind (kmph)	12	10	12	10	8	
200	250	90	90	Wind Direction	110	90	90	90	110	

## Weather Summary for Tiruppur district

## Weather forecast for next five days: (28.02.2024 to 03.03.2024)

According to district forecast issued by the India Meteorological Department for Tiruppur district, sky will be clear. Dry weather is expected on next 5 days. Maximum temperature is expected to be around 34°C to 36°C. Minimum temperature is expected to be around 23°C to 25°C. Morning relative humidity is expected to be around 80 per cent and evening relative humidity is expected to be around 20 per cent. Average wind speed is expected to be around 8-12 km per hour and the wind direction will be from South West direction.

Tiruppur District Block Wise Rainfall(mm) for next five days													
Date	Avinashi	Dharapuram	Gudimangalam	Kangeyam	Kundadam	Madathukualm	Mulanur	Palladam	Pongalur	Tiruppur	Udumalpet	Uthukuli	Vellakovil
28-02-2024	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
29-02-2024	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
01-03-2024	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
02-03-2024	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
03-03-2024	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

# Agro Advisory

Crop	Stage	Advisory
General		Dry days with mostly clear sky is expected during next week with wind
		speed of 6-12 km/hr in western zone districts. Both the maximum and
		minimum temperature are increasing while the relative humidity is
		decreasing.
		Irrigation may be done based on soil moisture and apply crop residue
		mulches to all irrigated crops to maintain sufficient soil moisture.
Rice	Flowering	Due to rising temperature (both maximum and minimum), sucking pest
		may appear in transplanted Navarai season rice. To control, spray
		Phosphamidon 40SL @ 600 ml / ha.
Sugarcane	Vegetative	Since higher temperature associated with low humidity is favourable
		for the early shoot borer incidence in sugarcane and hence farmers are
		requested to do frequent irrigation to keep the field under moist
		condition.
Banana		Bacterial hot rot disease is seen in some plantations of the 5 months old
		banana crop hence farmers are advised to monitor the crop and drench
		bleaching powder @ 20gm/liter of water.
		Provide support to banana above 5 months old as wind speed is around
		8 – 10 kmph.

Mango	Flowering	Considering the dry weather, if mango tree doesn't have flowers, Spray					
		either 0.5% urea (5g/litre) or 1% Potassium nitrate (10g/litre) to induce					
		flowering under sufficient soil moisture.					
		If mango has flowers, spray 2% Potassium nitrate to increase fruit set					
		and retardation of fruit, when sufficient soil moisture is available.					
Animal		Since the day temperature found increasing with decreasing relative					
Husbandry		humidity, provide enough clean water to both animals and poultry.					
		Hang wet gunnybags around the animal shed to reduce the					
		temperature. Do not allow animals for grazing during mid day.					

#### SMS advisory

> The day temperature is increasing, while the relative humidity is decreasing. Hence, provide enough clean water to both animals and poultry.

Professor and Head Principal Nodal Officer (GKMS) Agro Climate Research Centre