

Agromet Advisory Bulletin for the Nagaipattinam District

(From 03rd January 2024 to 07th January 2024)



Issued jointly by Tamil Nadu Rice Research Institute - Aduthurai and IMD

Bulletin no: 01/2024

Date: 02.01.2024

Weather Summary for the period since the issue of last bulletin no. 104 (29.12.2023) for Tamil Nadu: Nagaipattinam district

Maximum temperature of 30 °C, minimum temperature of 22 - 23 °C, isolated rainfall was recorded in Nagaipattinam district.

Weather Summary and forecast for Nagaipattinam district

Past week Summary 29.12.2023 to 01.01.2024				Parameters	Ensemble weather forecast valid until 08.30 hr of 07.01.2024				
Day-1	Day-2	Day-3	Day-4	Date	Day-1	Day-2	Day-3	Day-4	Day-5
29/12	30/12	31/12	01/01		03/01	04/01	05/01	06/01	07/01
4.7	0.3	0.0	0.0	Rainfall (mm)	3	2	2	2	26
30	30	30	30	Max. Temp. (°C)	29	29	28	28	28
24	26	25	23	Min. Temp. (°C)	23	23	23	22	22
5	4	5	4	Cloud cover (Octa)	8	8	7	6	8
95-80	80-76	81-76	83-71	RH I-II (%)	80-70	80-70	90-80	90-70	90-70
6	6	7	7	Wind (kmph)	14	14	12	12	14
North East	North East	North East	North East	Wind direction	North East	North East	North East	North East	East North East

Weather forecast for next five days (03.01.2024 to 07.01.2024)

According to district level forecast issued by India Meteorological Department for Nagaipattinam district, the sky will be moderate cloudy. Light rainfall is expected on 30.12.2023 to 01.01.2024. Maximum temperature is expected to be around 28 – 29 °C and minimum temperature is expected to be around 22 - 23 °C. Morning relative humidity is expected to be around 80 - 90 percent and evening relative humidity is expected to be around 70 - 80 percent. Average wind speed is expected to be around 12 – 14 km per hour. The wind direction will be north east.

Agro Advisory

Crop	Stage	Advisory		
Ger	neral	Farmers are requested to download and utilize the agricultural based mobile apps suggested by TNAU (TNAU AAS, Meghdoot and DAMINI).		
Rice	Main field	The pravelling weather condition favors leaf folder incidence. If leaf folder symptoms found (folded leaf, white scrape on leaf) spray Cartap hydrochloride 50 % SP @ 400 g/acre or Flubendiamide 20% WG 50 g/acre in 200 liter of water. The pravelling weather condition favors for Brown Plant Hopper incidence. To control this, Chlorantraniliprole 18.5% SC 60 g/ac or Clothianidin 50% WG 8-9.6 g/ac in 200 liter of water.		
		Sheath blight is being noticed in the paddy field due to prevailing weather condition. To control this, spray carbendazim 200g/ac or propiconozole 200 ml/acre in 200 liter water.		
Cattle	Cow	Due to cool night temperature, the livestock particularly the young ones should be kept under shelter with heating arrangements to prevent the incidence of pneumonia.		

SMS advisory

Farmers are requested to download and utilize the agricultural based mobile apps suggested by TNAU (TNAU AAS, Meghdoot and DAMINI).

Director TRRI, Aduthurai