Tamil Nadu Agricultural University



India Meteorological Department Agromet Advisory Bulletin for the Tiruppur District Bulletin No.037/2024 Issued on 07.05.2024 (From 08th May 2024 to 12th May 2024) Issued jointly by Agro Climate Research Centre, TNAU and IMD



Past Weather for Tamil Nadu District

Weather Summary for the period since the issue of last bulletin 036 (30.04.2024) for Tamil Nadu. Maximum temperature ranged from 36°C to 44°C and minimum temperature ranged from 25°C to 31°C, over the State. Kodaikanal and Uthagamandalam recorded maximum temperature ranged from 23°C to 29°C and minimum temperature ranged from 15°C to 16°C. During this period occurred rainfall was recorded in Tamil Nadu.

Tiruppur District Past Weather

In Tiruppur district, during 03.05.2024 to 06.05.2024 the Maximum temperature was recorded was 38°C to 39°C the minimum temperature was 26°C to 28°C. The morning relative humidity recorded was 66 per cent and evening relative humidity was 36 per cent. The average wind speed recorded was 9-23 km per hour and the wind direction were from East direction.

Past Week Summary (03.05.2024 to 06.05.2024)			Ensemble weather forecast valid until 08.30 hrs of 12.05.2024						
Day-1 03/04	Day-2 04/05	Day-3 05/05	Day-4 06/05	Date	Day-1 08/05	Day-2 09/05	Day-3 10/05	Day-4 11/05	Day-5 12/05
0.0	0.0	0.2	0.0	Rainfall	2	38	12	4	4
39	39	39	38	Max. Temp. (° C)	42	40	40	39	40
28	27	27	26	Mini. Temp. (° C)	26	24	23	23	22
7	7	5	4	Cloud cover (Octa)	5	8	6	7	7
61	66	65	51	RH morning (%)	70	70	70	70	70
42	46	44	36	RH evening (%)	30	30	30	40	30
18	9	23	21	Wind (kmph)	14	8	8	8	10
200	250	230	230	Wind Direction	250	230	250	160	250

Weather Summary for Tiruppur district

Weather forecast for next five days: (08.05.2024 to 12.05.2024)

According to district forecast issued by the India Meteorological Department for Tiruppur district, sky will be mostly. Moderate rainfall expected on next 5 days. Maximum temperature is expected to be around 39°C to 42°C. Minimum temperature is expected to be around 22°C to 26°C. Morning relative humidity is expected to be around 70 per cent and evening relative humidity is expected to be around 30 per cent. Average wind speed is expected to be around 8-14 km per hour and the wind direction will be from South West direction.

Tiruppur District Block Wise Rainfall(mm) for next five days													
Date	Avinashi	Dharapuram	Gudimangalam	Kangeyam	Kundadam	Madathukualm	Mulanur	Palladam	Pongalur	Tiruppur	Udumalpet	Uthukuli	Vellakovil
08-05-2024	0.0	0.0	0.0	0.0	0.0	1.0	0.0	0.0	0.0	0.0	1.0	0.0	0.0
09-05-2024	33.0	26.0	25.0	31.0	29.0	25.0	23.0	32.0	32.0	35.0	25.0	34.0	25.0
10-05-2024	0.0	1.0	1.0	0.0	0.0	2.0	1.0	0.0	0.0	0.0	2.0	0.0	0.0
11-05-2024	1.0	3.0	4.0	2.0	2.0	5.0	3.0	2.0	2.0	1.0	5.0	1.0	2.0
12-05-2024	1.0	3.0	4.0	1.0	2.0	5.0	2.0	2.0	1.0	1.0	5.0	1.0	1.0

Agro Advisory

Crop	Stage	Advisory
General		Rainfall is expected during this week, which will relive the heat stress
		during the later part of the week. The wind speed will be between 6 -
		14 km/hr.
		Irrigation may be done based on soil moisture during afternoon hours
		and apply crop residue mulches to all irrigated crops to maintain
		sufficient soil moisture.
		Plan for summer ploughing against the slope with soil inverting type
		ploughs.
		The saline concentration of irrigation water of wells and bore wells
		normally increases in summer months and this will create yellowing or
		browning of leaves leads to death of lower leaves. This symptom
		normally occurs in tapioca due to saline water irrigation. The following
		management measures may be adopted to reduce the damage.
		• Frequent irrigation to keep the soil moisture wet and to avoid
		accumulation of salt in the surface of the soil.
		• Irrigation may be done during early morning or late evening hours.
		• Application of organic manure to neutralize the salt.
		• Spraying of good quality water during evening and morning hours.
		 Mixing of good quality water with well water for irrigation.

Sugarcane	Vegetative	Considering the local circulation due to increased temperature and
		expected summer rains, provide propping to the sugarcane by tying
		opposite row to prevent from lodging.
Coconut		Summer rains during next two months may be harvested in root zone
		of coconut by deepening the tree basin with inward slopes. Till then,
		ensure to provide 100 liters of water per tree per day to maintain the
		productivity.
Animal		Increasing temperature and decreasing relative humidity may cause
Husbandry		discomfort to the animals. Hence, provide clean water to both animals
		and poultry.
		Application of water over the sheds of the cattle and poultry will
		reduce the heat stress or even hanging of wetted gunny bags will also
		reduce the temperature to a considerable extent.
		To avoid the heat stress, animals feed the animal during night or early
		morning or late evening hours.
		Summer rains may occur as thunderstorm during the afternoon hours.
		Safeguard the animals.

SMS advisory

Moderate rainfall with squally wind is expected. Hence farmers are requested to take suitable precautionary measures to avoid economical loss.

> Professor and Head Principal Nodal Officer (GKMS) Agro Climate Research Centre